



ACTIVE⁺

CYMOXI

SELENISED ORAL DRENCH FOR SHEEP

For the treatment of Gastrointestinal and Pulmonary Nematodes, and Itchmite of sheep.



ACTIVE +1

MOXIDECTIN

Active Constituent: 1mg/mL Moxidectin.
Contains 0.5mg/mL Selenium. Contains benzyl alcohol <50mg/mL.

USE ON:
SHEEP



TREATMENT OF:
GASTROINTESTINAL & PULMONARY NEMATODES,
ITCHMITE

CYMOXI

SELENISED ORAL DRENCH FOR SHEEP

Active Constituent: 1mg/mL Moxidectin. Contains 0.5mg/mL Selenium. Contains benzyl alcohol <50mg/mL.



BENEFITS

- Contains Moxidectin – a potent macrocyclic lactone (ML).
- Fast-acting broad-spectrum.
- Effective against *Haemonchus contortus* and *Ostertagia circumcincta*.
- Persistent activity up to 35 days.
- Provides a supplemental source of Selenium.
- Safe to use concurrently with other treatments.
- Developed in New Zealand for New Zealand.

ADMINISTRATION

SHAKE WELL BEFORE USE.

Invert drum three times before use. Administer 1mL per 5kg liveweight using any standard drenching equipment. Correct drenching technique should be used. Equipment should be checked for accuracy of dose delivery before drenching commences. Check accuracy of drench guns regularly.

DOSE RATE: 1mL/5kg liveweight to give 0.1mg/kg Selenium and 0.2mg/kg Moxidectin.

LIVE WEIGHT	DOSE VOLUME	LIVE WEIGHT	DOSE VOLUME
15 kg	3 mL	46 - 50 kg	10 mL
16 - 20 kg	4 mL	51 - 55 kg	11 mL
21 - 25 kg	5 mL	56 - 60 kg	12 mL
26 - 30 kg	6 mL	61 - 65 kg	13 mL
31 - 35 kg	7 mL	66 - 70 kg	14 mL
36 - 40 kg	8 mL	71 - 75 kg	15 mL
41 - 45 kg	9 mL	76 - 80 kg	16 mL

For animals over 80 kg use at a rate of 1 mL per 5 kg.

WITHHOLDING PERIODS

MEAT: Sheep producing meat and offal for human consumption must not be slaughtered during or within 10 days of the last treatment.

MILK: Milk intended for sale or human consumption must be discarded during treatment and for 35 days following the last treatment.

EFFICACY

Active+ Cymoxi Oral sheep drench contains Moxidectin; a member of the Milbemycin/Avermectin family. It is effective against sensitive strains of the following internal parasites;

Gastrointestinal Parasites:

Adult and immature stages of: *Haemonchus contortus*, *Teladorsagia circumcincta*, *Ostertagia trifurcata*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*, *Nematodirus spathiger*, *Nematodirus filicollis*, *Cooperia curticei* (adult only), *Cooperia oncophora* (adult only), *Cooperia punctata* (adult only), *Oesophagostomum venulosum* (adult only), *Chabertia ovina*, *Trichuris ovis* (adult only), *Strongyloides papillosus* (adult only), *Inhibited* (or hypobiotic) *larvae of Haemonchus, Ostertagia and Trichostrongylus.*

Lungworm (adult and immature):

Dictyocaulus filaria.

External Parasites:

Psorergates ovis (itchmite).

STORAGE

Store between 5°C and 25°C. Protect from light.

PACK SIZES

Active+ Cymoxi Oral is available in 20L, 10L and 5L packs.

Active+ Cymoxi Oral is registered pursuant to the ACVM Act 1997 No. A011500.

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited. PO Box 217234. Botany Junction. Auckland

