



ACTIVE⁺

TREBLE SHEEP ORAL

A TRIPLE COMBINATION DRENCH FOR SHEEP

For the treatment and control of internal parasites in sheep.



ACTIVE +3

Active Ingredients: 1g/L abamectin, 22.7g/L oxfendazole, 40.0g/L levamisole hydrochloride, 2.5g/L cobalt, 0.5g/L selenium.

**ABAMECTIN
OXFENDAZOLE
LEVAMISOLE**

USE ON:
SHEEP 

MANAGEMENT OF:
INTERNAL PARASITES

TREBLE SHEEP ORAL

A TRIPLE COMBINATION DRENCH FOR SHEEP

Active Ingredients: 1g/L abamectin, 22.7g/L oxfendazole, 40.0g/L levamisole hydrochloride, 2.5g/L cobalt, 0.5g/L selenium



BENEFITS

- Convenient 1ml/5kg dose rate.
- For the management of internal parasites in sheep.
- White/ML/Clear combination plus Selenium and Cobalt.
- Aids in the supplementation of Selenium and Cobalt.
- Cost effective.
- Developed in New Zealand for New Zealand.

ADMINISTRATION

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

SHAKE WELL BEFORE USE.

Do not use in lambs under 20kg or under 4 weeks of age.

A representative sample of animals should be weighed before treatment.

Dose rate to be based on the heaviest animal in each group (see precautions). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Regularly check the accuracy of the drench gun.

Correct drenching technique should be used.

DOSE RATE 1mL/5kg liveweight to give 0.2mg/kg abamectin, 4.5mg/kg oxfendazole, 8.0 mg/kg levamisole HCL, 0.1mg/kg selenium and 0.5mg/kg cobalt.

Active+ Treble Sheep Oral is to be administered via the oral route.

| DOSAGE RATE 1mL/5kg Liveweight | | |
|--------------------------------|------------|--------------------|
| LIVE WEIGHT | DOSE | DOSES PER 20L PACK |
| Less than 20 kg | Do not use | |
| 20 kg | 4 mL | 5000 |
| 30 kg | 6 mL | 3333 |
| 40 kg | 8 mL | 2500 |
| 50 kg | 10 mL | 2000 |
| 60 kg | 12 mL | 1666 |
| 70 kg | 14 mL | 1428 |
| 80 kg | 16 mL | 1250 |
| Over 80 kg | 1 mL/ 5kg | - |

WITHHOLDING PERIODS

MEAT: 14 days - Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of the last treatment.

MILK: 35 days - Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

STORAGE

Store in a cool dry place at room temperature (i.e. 5°C to 25°C).

PACK SIZES

Active+ Treble Sheep Oral is available in 10L, and 20L packs.

EFFICACY

Active+ Treble Sheep Oral contains abamectin, a member of the macrocyclic lactone (ML) drench family levamisole hydrochloride, a member of the levamisole (clear) drench family and oxfendazole a member of the benzimidazole (white) drench family. Also contains chelated cobalt and sodium selenate.

Anthelmintic resistant strains: Active+ Treble Sheep Oral Drench may be effective in treating some populations of gastrointestinal worms with single or dual resistance to the avermectin/milbemycin, benzimidazole or levamisole/morantel anthelmintic action families. The resistant populations must be susceptible to one or more of the anthelmintic actives in the drench for it to be effective.

Aids in selenium and cobalt supplementation.

At the recommended dose, this product is effective for the treatment and control of sensitive strains of the following mature and immature parasites of sheep (* includes L4 inhibited larvae.).

| COMMON NAME | GENUS | SHEEP SPECIES |
|-----------------------------|-------------------------|---|
| ROUNDWORMS | | |
| ABOMASUM | | |
| Barber's Pole Worm* | <i>Haemonchus</i> | <i>contortus</i> |
| Small Brown Stomach Worm* | <i>Teladorsagia</i> | <i>circumcincta</i> |
| Stomach Hair Worm | <i>Trichostrongylus</i> | <i>axei</i> |
| SMALL INTESTINE | | |
| Black Scour Worm | <i>Trichostrongylus</i> | <i>colubriformis</i> <i>vitrinus</i> |
| Small Intestinal Worm* | <i>Cooperia</i> | <i>curticei</i> |
| Thin-Necked Intestinal Worm | <i>Nematodirus</i> | <i>filicollis</i> <i>spathiger</i> |
| Hookworm | <i>Strongyloides</i> | <i>papillosus</i> |
| LARGE INTESTINE | | |
| Large Bowel Worm | <i>Oesophagostomum</i> | <i>venulosum</i> |
| Large-Mouthed Bowel Worm | <i>Chabertia</i> | <i>ovina</i> |
| Whipworm | <i>Trichuris</i> | <i>ovis</i> |
| LUNGWORM | | |
| Large Lungworm | <i>Dictyocaulus</i> | <i>filaria</i> |

PRECAUTIONS

Doses of three or more times those recommended can cause symptoms of levamisole toxicity, so estimate liveweights carefully.

Animals under stress due to advanced pregnancy, adverse weather conditions, poor nutrition, prolonged yarding or dehydration may be more susceptible to toxicity.

Fatal interactions may occur between levamisole and organophosphate dips.

Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian.

Registered pursuant to the ACVM Act 1997, No. A011892. See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR100758. See www.epa.govt.nz for controls

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland