


Animec Super



SUPERIOR STRENGTH, PROTECTION & CHOICE

Animec Super

Solution for injection for Cattle

Contains 1% w/v Ivermectin and 10% w/v Clorsulon

One dose solution for the treatment and control of parasites, worms and liver fluke.

Colourless to pale yellow non-aqueous solution for injection in cattle. Contains ivermectin and the flukicide clorsulon. Its convenience, broad spectrum efficacy and wide safety margin make it an ideal product for parasite control in cattle.



New Zealand owned and operated

Manufactured for and distributed by AHD Ltd

1229 Maraekakaho Road Hastings

Phone: 06 873 3611. Email: sales@ahdltd.co.nz

www.ahdltd.co.nz

Animec Super

Animec Super is effective for the treatment and control of the following in cattle:

- Gastro-intestinal roundworms
- Lungworms
- Eye worms (adult)
- Warbles (parasitic stages)
- Mange mites
- Sucking lice
- Adult liver fluke

Effects of parasites

- Loss of body condition
- Weight loss
- Vulnerable to other diseases
- Damage to stomach lining
- Lower feed absorption
- Reduction in reproductive performance
- Coughing
- Scouring

When to dose?

1. Establish a dosing programme with your veterinary surgeon.
2. It is recommended dosing programmes commence in the early autumn (February to May) period and again August/September.
3. It is also recommended to dose the cattle prior to entering feedlots or housing.
4. The interval between doses will depend on the level of pasture contamination, weather conditions and geographical location.

Dosage

| WEIGHT (kg) | DOSE Volume ml | DOSES 500ml pk | WEIGHT (kg) | DOSE Volume ml | DOSES 500ml pk |
|-------------|----------------|----------------|-------------|----------------|----------------|
| 50-100 | 2 | 250 | 351-400 | 8 | 62 |
| 101-150 | 3 | 166 | 401-450 | 9 | 55 |
| 151-200 | 4 | 125 | 451-500 | 10 | 50 |
| 201-250 | 5 | 100 | 501-550 | 11 | 45 |
| 251-300 | 6 | 83 | 551-600 | 12 | 41 |
| 301-350 | 7 | 71 | 601-650 | 13 | 38 |

Consult your veterinary surgeon to establish a dosing schedule. Know the weight of the animal before administering.

Not permitted for use in animals producing milk for human consumption, including pregnant animals intended to produce milk for human consumption.

Withholding cattle and bobby calves: 28 days. Milk 14 days.

