Product Information



Name:	ANTHEM® SHEEP HI MIN			
Pack Sizes/Codes:	20L / DAHM20	CONTROL OF AN AND AND AND AND AND AND AND AND AND		
Main Points:	 A double action combination drench containing both levamisole and albendazole which is a member of the benzimidazole family. Combines two anthelmintics with alternative modes of action Prevention of selenium deficiencies Includes minerals and selenium Standard 12ml/5kg sheep dose rate Control of BZ and levamisole resistant parasites 			
Status:	OTC			
Ingredients:	 37.5g/L levamisole 23.8g/L albendazole HCI Each 10mL dose contains: 5mg selenium 2.5mg cobalt 21mg copper 			
Indications:	 For the control of sensitive mature and immature Haemonchus, Ostertagia, Trichostrongylus, Nematodirus, Cooperia, Strongyloides, Bunostomum, Oesophagostomum, Chabertia, Trichuris, Dictyocaulus, Moniezia and adult liver fluke in sheep. Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment. 			
Action:	A double action combination drench containing both levamisole and albendazole which is a member of the benzimidazole family. Highly effective against internal parasites sensitive to one or both of these families.			
	By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.			
	SHAKE WELL BEFORE USE			
Administration:	ANTHEM® SHEEP HI MIN is a ready to use combination drench and can be given using standard drenching equipment. Drench guns should be rinsed with water immediately after use. Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment. Dose rate to be based on the heaviest animal in each group. Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label direction for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching technique should be used. Check the accuracy of drench guns regularly.			



Animal Health Direct Limited PO Box 8015 | Havelock North | 4157 | t: 06 873 3611 e: sales@ahdltd.co.nz | ahdltd.co.nz



Product Information

|--|

	Withholding periods –		
	Milk: Milk intended for human consumption or manufacture for human consumption must be discarded during treatment and for 35 days following the last treatment.		
	Meat: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of the last treatment.		
	Dose at 1mL per 5kg which gives 7.5 mg/kg levamisole HCl with 4.75mg/kg albendazole.BodyweightDose20L Treats		
Dosage:	15kg 16-20kg	3mL 4mL	6666 5000
	21-25kg 26-30kg 31-35kg	5mL 6mL 7mL	4000 3333 2857
	36-40kg 41-45kg 46-50kg Above 50 kg add	8mL 9mL 10mL d 1 mL per 5 k	2500 2222 2000 a bodyweight
	May be harmful if swallowed. Causes mild skin irritation and may cause an allergic skin reaction. Suspected of causing genetic defects. May damage fertility or the unborn child. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. If medical advice is needed, have product container or label at hand.		
Special Precautions:	Handling Precautions - Avoid skin contact. Wear protective gloves. Do not breathe vapours/spray. Wash hands thoroughly after using. Wash contaminated clothing before reuse.		
	First Aid - If on skin wash with plenty of soap and water. If skin irritation or rash develops or if you feel unwell contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.		
	Environmental Protection - Very toxic to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment. Avoid release to the environment. Collect spillage.		
Storage:	Store locked up in a cool place with lid on.		
	Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse empty container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.		
Registration:	Registered pursuant to the ACVM Act 1997, No. A011153		
Category:	Parasite control drench		



