


Name:	<b>Bovaid</b>	
Pack Size/s	10L	
AHD Product Code:	AHDBACO10	
Main Points:	A combination drench for the control of roundworms, tapeworm, lungworm and adult liver fluke in cattle including roundworms resistant to benzimidazole or levamisole	
Status:	OTC/VET	
Ingredients:	<ul style="list-style-type: none"><li>• 100 g/L Albendazole</li><li>• 75 g/L Levamisole HCl</li></ul>	
Indications:	<ul style="list-style-type: none"><li>• BOVAID® COMBINATION DRENCH FOR CATTLE is a combination anthelmintic containing a member of both benzimidazole (white) and imidazothiazole (clear) drench families.</li><li>• It is effective against parasites sensitive to one or both of these families.</li></ul>	
Action:	<ul style="list-style-type: none"><li>• For the control of sensitive mature and immature Haemonchus, Ostertagia, Trichostrongylus, Nematodirus, Cooperia, Triclyoides, Bunostomum, Oesophagostomum, Chabertia and Trichuris that may be resistant to benzimidazole or levamisole.</li><li>• Also for the control of tapeworm, lungworm and adult liver fluke in cattle.</li></ul>	
Administration:	<ul style="list-style-type: none"><li>• Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment. Dose rate to be based on the heaviest animal in each group.</li><li>• Do not under dose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range.</li><li>• Drafting into two or more lines may be appropriate, to avoid excessive overdosing.</li><li>• Correct drenching techniques should be used. Check the accuracy of drench guns regularly.</li></ul>	
Dosage:	Cattle 10mg/kg Albendazole and 7.5mg/kg Levamisole HCl using 1ml/10kg live weight.	
Special Precautions:	<p>Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.</p> <p>Doses of 3 or more times recommended dose can cause symptoms of levamisole toxicity, so estimate liveweights carefully.</p>	



# Product Information

**Bovaid**

	<p>Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.</p> <p><b>WITHHOLDING PERIODS:</b></p> <p><b>Meat</b> – Animals producing meat of offal for human consumption must not be sold for slaughter during or within 14 days of cessation of last treatment.</p> <p><b>Milk</b> – Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.</p>
Storage:	<p>Store in a cool place below 30°C. Store locked up.</p> <p>KEEP OUT OF REACH OF CHILDREN.</p>
Registration:	<p>Registered pursuant to the ACVM Act 1997 No A11145.</p> <p>See <a href="http://www.foodsafety.govt.nz">www.foodsafety.govt.nz</a> for registration conditions</p>
Category:	<p>Parasitic control</p>

**Animal Health Direct Limited**



PO Box 8015 | Havelock North | 4157 | t: 06 873 3611  
e: [sales@ahdltd.co.nz](mailto:sales@ahdltd.co.nz) | [ahdltd.co.nz](http://ahdltd.co.nz)

