

DANGER
KEEP OUT OF REACH OF CHILDREN
For Animal Treatment Only



Supporting Kiwi Farmers



COMBINATION CATTLE DRENCH

A broad spectrum oral combination (containing albendazole and levamisole) cattle drench for the control of susceptible gastrointestinal roundworms and lungworms in cattle including benzimidazole or levamisole resistant strains. Also for the control of tapeworms and adult liver fluke. Also contains minerals to provide supplementary nutritional support.

INGREDIENTS:

Albendazole 100 mg/mL, Levamisole HCl 75 mg/mL, Selenium (as sodium selenate) 0.9 mg/mL 2.2 mg/mL, Cobalt (as chelate) 0.2 mg/mL, Copper (as chelate) 2 mg/mL Zinc (as chelate) 0.6 mg/mL, Iodine (as iodate) 1 mg/mL

NET CONTENTS

5 Litre

LCCD5/07



180mm deep

COMBINATION CATTLE DRENCH

PLEASE READ LABEL COMPLETELY BEFORE USE



Precautions: Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose or dose more frequently than three weekly. Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity so estimate live weights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.

Handling Precautions: Do not handle until all safety precautions have been read and understood. Harmful if swallowed. May cause an allergic skin reaction. Albendazole and levamisole hydrochloride are suspected of causing genetic defects and/or may damage fertility or the unborn child. Levamisole hydrochloride may possibly affect blood and the haemopoietic system through prolonged or repeated exposure. Wash hands and exposed skin before meals and after use. Do not eat, drink, or smoke when using this product. Do not breathe fumes, mist, vapours, or spray. Avoid skin contact. Wear protective gloves/protective clothing and use personal protective equipment as required. Contaminated work clothing should not be allowed out of the workplace. Take off and wash contaminated clothing before reuse. **First Aid:** If swallowed, rinse mouth. Do NOT induce vomiting. For advice, or if you feel unwell, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately. If on skin, wash with plenty of soap and water. If skin irritation or rash occurs, or if exposed or concerned or you feel unwell, get medical advice/attention.

ENVIRONMENTAL PROTECTION Hazardous to terrestrial vertebrates. Avoid release to the environment.

Emergency Response: In case of spillage wear appropriate clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

Disposal: Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

INDICATIONS

Contains albendazole (100 mg/mL), levamisole HCl 75 mg/mL, selenium (as sodium selenate 0.9 mg/mL, cobalt (as chelate) 0.2 mg/mL, copper (as chelate) 2 mg/mL, zinc (as chelate) 0.6 mg/mL and iodine (as iodate) 1 mg/mL. For the treatment of the following susceptible adult and immature stages of gastrointestinal roundworms and lungworms, including benzimidazole or levamisole resistant strains: barber's pole worm (*Haemonchus sp*), brown stomach worm (*Ostertagia spp*), stomach hair worm and black scour worm (*Trichostrongylus spp*), thread-necked intestinal worm (*Nematodirus spp*), small intestinal worm (*Cooperia spp*), intestinal threadworm (*Strongyloides sp*), hookworm (*Bunostomum sp*), (large-mouthed bowel worm (*Chabertia sp*), nodule worm (*Oesophagostomum sp*), (whipworm (*Trichuris sp*), Lungworm (*Dictyocaulus spp*). Also, for the control of tapeworm (*Moniezia sp*) and liver fluke (*Fasciola hepatica* -adults only). Also kills roundworm and fluke eggs.

RESISTANCE MANAGEMENT:

Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinary practitioner for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DIRECTIONS FOR USE: Shake well before use. For oral dosing only. Combination Cattle Drench is a ready to use oral anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment, weigh a representative sample of Cattle to determine the weight of the heaviest animal. Where there is a large variation in liveweights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band. Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

DOSE CHART

Oral dose rate cattle: 1mL/10kg (equivalent to 10 mg/kg albendazole, 7.5 mg/kg levamisole HCl, 0.09 mg/kg selenium, 0.02 mg/kg cobalt, 0.2 mg/kg copper, 0.06 mg/kg zinc and 0.1 mg/kg iodine).

ANIMAL WEIGHT	DOSE	NO. OF DOSES PER 5L PACK
60 kg	6 mL	833
100 kg	10 mL	500
150 kg	15 mL	333
200 kg	20 mL	250
250 kg	25 mL	200
300 kg	30 mL	166
350 kg	35 mL	142
400 kg	40 mL	125
450 kg	45 mL	111
500 kg	50 mL	100
550 kg	55 mL	90
600 kg	60 mL	83

Over 600kg dose at the rate of 1mL/10 kg.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

MEAT: Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within **10 days** of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than **35 days** following the last treatment.

STORAGE: Store locked up in original container, tightly closed, in cool, dry conditions.

Protect from light. Avoid temperatures below 0° and above 25°C.

Registered Pursuant to the ACVM Act 1997, No. A9904.

See www.foodsafety.govt.nz for registration conditions.

Registered to: Seneca Holdings Limited

Distributed by: Animal Health Direct Ltd.

2 Tumu Way, Longlands, Hastings, New Zealand.

sales@ahd ltd.co.nz. Ph (06) 873 3611.

Batch No.:

Expiry date:



9 417726 702636

LCCDS5B/07