






# COXI-STOP™ 50 Coccidiocide for Piglets and Cattle

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL

## Low Volume, High Impact - Coccidiosis Control Made Simple.

For the control of coccidiosis in neonatal piglets caused by *Isoospora suis*, and in cattle up to 9 months of age caused by *Eimeria bovis* or *Eimeria zuernii*.

-  Easy to use suspension
-  Low volume 1mL dose for piglets
-  Contains the active Toltrazuril
-  Can be used in piglets from 3-6 days of age
-  Can be used in calves up to 9 months of age



	COXI-STOP™ 50	BAYCOX® Piglet & Cattle
Active Constituent	Toltrazuril 50g/L	Toltrazuril 50g/L
Species	Cattle	Cattle
Route of Administration	Orally	Orally
Does rate	Piglets: 1mL Calves: 3mL per 10kg	Piglets: 1mL Calves: 3mL per 10kg
<b>Withholding Periods</b>		
Pigs: Meat	70 days	70 days
Calves: Meat	56 days	56 days
Cattle: Milk	DO NOT USE	DO NOT USE
ESI: Cattle	75 days	75 days

COXI-STOP™ is a trademark of Abbey Laboratories Pty Ltd.

BAYCOX® is a registered trademark of Bayer Intellectual Property GmbH.

# COXI-STOP™ 50 Coccidiocide for Piglets and Cattle

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL



Supporting Kiwi Farmers

**For the control of coccidiosis in neonatal piglets caused by *Isoospora suis*,  
and in cattle up to 9 months of age caused by *Eimeria bovis* or *Eimeria zuernii*.**

## INDICATIONS

COXI-STOP™ 50 is effective in one dose because it attacks all stages of the parasite in the animal.

**Piglets:** COXI-STOP™ 50 controls coccidiosis caused by *Isoospora suis* infection of young piglets. *Isoospora suis* infection is a major cause of diarrhoea, dehydration, and occasional mortality in piglets between birth and 3-4 weeks of age.

**Cattle:** COXI-STOP™ 50 is for the treatment of clinical coccidiosis, and as an aid in the prevention of coccidiosis in cattle up to 9 months of age. COXI-STOP™ 50 controls coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in young cattle. Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood strained. Recently weaned calves are susceptible to outbreaks of clinical disease.

## DIRECTIONS FOR USE

### Restraints

**DO NOT RE-TREAT** calves for 75 days after the last treatment.

**DO NOT TREAT** piglets later than 8 days of age.

## DOSAGE AND ADMINISTRATION

### SHAKE THE CONTAINER BEFORE USE.

The product is administered orally. Use the measuring cup to give the full dose by mouth.

## DOSE RATE TABLE

	Dose	Prevention	Treatment
<b>Piglets</b>	1mL	Treat at 3-6 days old	Treat at onset of coccidial diarrhoea. Treat all piglets in infected litters and all infected and 'at risk' litters in the farrowing house.
<b>Cattle</b>	3mL per 10kg bodyweight	In the case of history of coccidiosis on farm, treat 1 week before expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves, treat at weaning time when milk feeding ceases.	Treat at onset of diagnosed clinical signs.

## GENERAL DIRECTIONS

**Piglets:** In the case of diarrhoea in neonatal piglets, obtain a veterinary diagnosis of coccidiosis before using this product. Clinical coccidiosis appears in piglets from 5 to 30 days of age. Control of the disease is most effective when COXI-STOP™ 50 is given as a preventive before intestinal damage occurs and diarrhoea begins.

**Cattle:** In the case of diarrhoea in calves, obtain a veterinary diagnosis of coccidiosis before using this product. Early treatment of calves as soon as the diagnosis is obtained will prevent further intestinal damage in affected animals and prevent infection of the herd. To obtain maximum benefit on farms with a history of coccidiosis, COXI-STOP™ 50 should be given approximately 1 week before expected onset of clinical signs. Treatment of newly weaned calves at the time of milk withdrawal will control coccidiosis associated with weaning.

## WITHHOLDING PERIODS

**MEAT (PIGS): DO NOT USE less than 70 days before slaughter for human consumption.**

**MEAT (CATTLE): DO NOT USE less than 56 days before slaughter for human consumption.**

**MILK (CATTLE): DO NOT USE in lactating or pregnant cows where milk may be used or processed for human consumption.**

## TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI) PIGS: An ESI has not been established for this product for pigs. Note- observing the meat withholding period may not be sufficient**

**to mitigate potential risks to export trade. Trade advice should be sought from Abbey Animal Health Pty Ltd on 02 8088 0720 before using this product.**

## EXPORT SLAUGHTER INTERVAL (ESI)

**CATTLE: DO NOT slaughter cattle for export for 75 days after last treatment. The ESI for cattle on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 02 8088 0720 or the APVMA website: ([www.apvma.gov.au/residues/](http://www.apvma.gov.au/residues/))**

## SAFETY DIRECTIONS

When using this product wear rubber gloves. After each day's use wash gloves.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

**ENVIRONMENTAL PROTECTION:** Do not contaminate ponds, waterways or drains with this product or used container.

## DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## STORAGE

Store below 30°C (room temperature). Protect from light.

COXI-STOP™ is a trademark of Abbey Laboratories Pty Ltd.

Registered pursuant to the ACVM Act 1997, No A11753.