








COXI-STOP™ 50 COCCIDIOCIDES FOR PIGLETS AND CATTLE

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL

Low Volume, High Impact - Coccidiosis Control Made Simple.

For the control of coccidiosis in neonatal piglets caused by *Isospora suis*, and in cattle up to 9 months of age caused by *Eimeria bovis* or *Eimeria zuernii*.

-  Easy to use suspension
-  Low volume 1mL dose for piglets
-  Contains the active Toltrazuril
-  Can be used in piglets from 3-6 days of age
-  Can be used in calves up to 9 months of age



	COXI-STOP™ 50	BAYCOX® Piglet & Cattle
Active Constituent	50g/L TOLTRAZURIL	50g/L TOLTRAZURIL
Species	Cattle	Cattle
Route of Administration	Orally	Orally
Dose rate	Piglets: 1mL Calves: 3mL per 10kg	Piglets: 1mL Calves: 3mL per 10kg
Withholding Periods		
Pigs: Meat	49 days	70 days
Cattle: Meat	56 days	56 days

COXI-STOP™ is a trademark of Abbey Laboratories Pty Ltd.
Registered pursuant to the ACVM Act 1997, No A11753.

BAYCOX® is a registered trademark of Bayer Intellectual Property GmbH Ltd. APVMA Number 68946/130682.

COXI-STOP™ 50 COCCIDIOCID FOR PIGLETS AND CATTLE

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL



Supporting Kiwi Farmers

For the control of coccidiosis in neonatal piglets caused by *Isospora suis*, and in cattle up to 9 months of age caused by *Eimeria bovis* or *Eimeria zuernii*.

INDICATIONS

PIGLETS: COXI-STOP™ 50 controls coccidiosis caused by *Isospora suis* infection of young piglets. *Isospora suis* infection is major cause of diarrhoea, dehydration, loss of appetite, poor growth, and occasional mortality in piglets between birth and 3-4 weeks of age.

CATTLE: COXI-STOP™ 50 controls coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in young cattle. Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood stained. Recently weaned calves are susceptible to outbreaks of clinical disease. COXI-STOP™ 50 is effective in one dose because it attacks all stages of the parasite in the animal.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Thoroughly shake container before use.

PIGLETS

Give 1mL of COXI-STOP™ 50 COCCIDIOCID FOR PIGLETS AND CATTLE per piglet by mouth. Use once when the piglets are 3-6 days old, or as soon as coccidial diarrhoea occurs. Treat all piglets in infected litters and all infected and at-risk litters in the farrowing house.

CATTLE

Give 3mL of COXI-STOP™ 50 COCCIDIOCID FOR PIGLETS AND CATTLE per 10kg of bodyweight by mouth. For the treatment of clinical disease treat all affected and in contact animals. For prevention of coccidiosis on farms with a known history treat prior to the expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves treat at weaning time when meal feeding ceases.

GENERAL INSTRUCTIONS

Before using this product obtain a veterinary diagnosis for diarrhoea.

To achieve effective control of Coccidiosis good management and hygiene are vital. In animals already showing signs of diarrhoea, additional supportive therapy may be required.

Seek advice from a veterinarian or animal health advisor regarding effective management and control of Coccidiosis.

PIGLETS

Clinical coccidiosis of piglets appears in piglets from 5 to 30 days of age. Control of the disease is most effective when COXI-STOP™ 50 COCCIDIOCID FOR PIGLETS AND CATTLE is given as a preventative before intestinal damage occurs and diarrhoea begins. Therefore it is important to establish the correct cause of scouring in neonatal piglets then treat all new litters at 3-6 days of age to control the disease.

CATTLE

Early treatment of calves as soon as clinical signs of disease are seen and confirmed to be due to coccidiosis will reduce the impact of an outbreak by preventing further intestinal damage in affected animals and preventing non-affected animals from developing diarrhoea. To obtain maximum benefit on farms with a history of coccidiosis, COXI-STOP™ 50 needs to be given approximately 1 week prior to the expected onset of clinical signs. Treatment of newly weaned calves at the time of meal withdrawal will control coccidiosis associated with weaning.

DOSE RATE TABLE

	Dose	Prevention	Treatment
Piglets	1mL	Treat at 3-6 days old	Treat at onset of coccidial diarrhoea. Treat all piglets in infected litters and all infected and 'at risk' litters in the farrowing house.
Cattle	3mL per 10kg bodyweight	In the case of history of coccidiosis on farm, treat 1 week before expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves, treat at weaning time when meal feeding ceases.	Treat at onset of diagnosed clinical signs.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Pigs producing meat and offal for human consumption must not be sold for slaughter during or within 49 days of the last treatment.

Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 56 days of the last treatment.

HANDLING PRECAUTIONS

Toltrazuril possibly may affect development. Avoid contact with skin.

FIRST AID

If on skin wash off with soap and water. If you feel unwell seek medical advice. Call 0800 POISON (0800 764 766) or a doctor immediately.

ENVIRONMENTAL PROTECTION

Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

DISPOSAL

Preferably dispose of product by use. Triple rinse empty container, puncture or crush and dispose of at an approved landfill/ facility or if appropriate recycle. Otherwise dispose of product at an approved landfill or other approved facility.

STORAGE

Store below 30°C (room temperature).

Registered pursuant to the ACVM Act 1997, No A11753. See www.foodsafety.govt.nz for registration conditions.

COXI-STOP™ is a trademark of Abbey Laboratories Pty Ltd