

**WARNING**  
**KEEP OUT OF REACH OF CHILDREN**  
**FOR ANIMAL USE ONLY**

# DETONATE WORMER



## Parasite control for the health of your horse

Detonate for Horses effectively controls gastrointestinal roundworms, including the arterial stages of *Strongylus vulgaris*, small strongyles (including benzimidazole resistant strains), lungworm and bots in horses.

### ACTIVE INGREDIENTS

IVERMECTIN 4mg/mL

### DOSAGE

One syringe division of gel per 100kg bodyweight.  
Each syringe is sufficient to treat a 600kg horse.  
See reverse of label for full details.

### WITHHOLDING PERIOD

MEAT: 63 days

**STORAGE:** Protect from direct sunlight. Store below 25°C.

Registered pursuant to the ACVM Act 1997, No. A9607  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

REGISTERED TO SENECA HOLDINGS LTD

DISTRIBUTED BY:

AHD LTD

2 TUMU WAY, LONGLANDS, HASTINGS

EMAIL: [SALES@AHDLTD.CO.NZ](mailto:SALES@AHDLTD.CO.NZ)



*Supporting Kiwi Horses*



DET060325

PEEL HERE

Batch:

**30ML NET**

# DETONATE WORMER

Controls gastrointestinal roundworms, including the arterial stages of *Strongylus vulgaris*, small strongyles (including benzimidazole resistant strains), lungworm and bots in horses. This easy to administer gel is safe to use in horses of all ages including lactating mares or at any stage of pregnancy. It will not adversely affect the fertility of stallions.

## INDICATIONS:

Detonate for Horses kills gastrointestinal roundworms, including the arterial stages of *Strongylus vulgaris*, small strongyles (including benzimidazole resistant strains), lungworm and bots in horses. Detonate for Horses, given at the recommended dose rate, is indicated for the treatment and control of the following internal parasites of horses: **Large strongyles (redworms):** adults and arterial larval stages of *Strongylus vulgaris*, adults and tissue larval stages of *S. edentatus*, adults of *S. equinus* and *Triodontophorus* spp. Adult and immature small strongyles (redworms): including benzimidazole resistant strains: *Cyathostomum* spp., *Cylicocycylus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp., *Gyalocephalus* spp. Adult and immature lungworms: *Dictyocaulus arnfieldi* Pinworms: Adult and immature *Oxyuris equi* Ascarids: Adult and immature *Parascaris equorum* Stomach worms: Adult *Trichostrongylus axei*, *Habronema muscae* Intestinal threadworms: *Strongyloides westeri* Oral and gastric stages of bots: *Gasterophilus* spp.

## DOSAGE:

Detonate for Horses is to be dosed at the rate of one syringe division of gel per 100kg bodyweight, or the contents of one syringe to a 600kg horse. This provides 200µg ivermectin/per kilogram bodyweight.

By law, the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress to the treated animal when using the product in a manner other than as recommended on the label.

## ADMINISTRATION:

Remove the cap from the syringe tip. Insert the syringe tip between the lips at the interdental space (the gap between the front and back teeth). Depress the plunger and deposit the paste on the top of the tongue toward the back of the mouth. The horse's head should be raised for a few seconds after dosing. For best results all horses in a yard or grazing together should be included in a regular parasite control programme and treated at the same time. Foals should initially be treated from 6-8 weeks of age. Consult your veterinarian to develop a programme that meets your needs.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

## WITHHOLDING PERIOD:

**MEAT:** Horses producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 63 days of the last treatment.

## PRECAUTIONS:

Detonate for Horses has been formulated for use in horses only. It should not be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

## WARNING

May cause an allergic skin reaction. May cause harm to breast-fed children.

## HANDLING PRECAUTIONS

Do not handle until all safety precautions have been read and understood. Avoid contact during pregnancy and while nursing. Do not breathe fumes/mist/vapours/spray. Avoid skin contact. Wear protective gloves. Do not eat, drink or smoke when using this product. Wash hands and exposed skin before meals and after use. Contaminated clothing should not be allowed out of the workplace. Take off contaminated clothing and wash before reuse.

## FIRST AID:

If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If on skin wash with plenty of water. If skin irritation or rash occurs, get medical advice/attention. If exposed or concerned, get medical advice/attention.

## ENVIRONMENTAL PROTECTION:

Very toxic to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Hazardous to terrestrial invertebrates. Hazardous to soil organisms. Avoid release to the environment. Collect spillage.

## DISPOSAL:

Do not contaminate surface water or ditches with product or used containers. Containers and any residual contents should be disposed of safely, i.e. putting in rubbish or by burying in waste ground away from watercourses.

## STORAGE:

Protect from direct sunlight. Store below 25°C.

