

# AHD LTD PARALOCK™

## ABAMECTIN POUR ON FOR CATTLE

For the treatment and control of internal  
and external parasites of cattle.

## NEW PRODUCT RELEASE



## PRODUCT USAGE

Paralock Abamectin Pour-on for Cattle, when used at the recommended dose level of 500 micrograms abamectin per kg live weight, effectively controls the following parasites of cattle:

- **Gastrointestinal roundworms:** adult and immature *Ostertagia* spp (including inhibited immatures), *Haemonchus* spp, *Trichostrongylus* spp, *Cooperia* spp, *Bunostomum* spp, *Oesophagostomum* spp, and adults of *Chabertia* spp, *Nematodirus* spp, *Trichuris* spp, and *Strongyloides* spp.
- **Lungworms:** adult and immature *Dictyocaulus viviparus*.
- **Biting and Sucking Lice:** *Bovicola bovis*, *Linognathus vituli*.

Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Check accuracy of pour-on applicator regularly.

## PRODUCT BENEFITS

- 🐮 Meat withholding  
Cattle 35 days.  
Milk 35 days.
- 🐮 Effectively controls  
parasites in cattle.
- 🐮 Ready to use rainfast  
pour-on.



## PARALOCK INDICATIONS:

A representative sample of animals should be weighed before treatment. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc.). Do not under-dose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Paralock Abamectin Pour-On is rainfast, even if rain falls 2 hours or more after treatment.

### Contraindication

Do not use this product in calves under 4 months of age and under 100 kg liveweight.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

### Withholding periods

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 35 days of the last treatment.

Storage: Store locked up below 25°C. Protect from sunlight and keep it well closed.

## DOSE RATES:

The dose rate for cattle is 500 µg abamectin per kg bodyweight, which is provided by 1 mL Paralock Abamectin Pour-On per 20 kg bodyweight, or according to the table.

Weight range	Dose	Doses per 10 L
101-150 kg	7.5 mL	1333
151-200 kg	10 mL	1000
201-250 kg	12.5 mL	800
251-300 kg	15 mL	666
301-350 kg	17.5 mL	571
351-400 kg	20 mL	500
401-450 kg	22.5 mL	444
451-500kg	25 mL	400
501-550 kg	27.5 mL	363
551-600 kg	30 mL	333
601-650 kg	32.5 mL	307

Heavier animals (e.g. mature bulls) should receive an additional 1 mL for each 20 kg over 650 kg. Drum tap located behind back label.

## ACTIVE INGREDIENTS

Abamectin 10 mg/mL

Contains >941 g/L diethylene glycol monobutyl ether.

## CONTACT US

Animal Health Direct Ltd  
2 Tumu Way, Longlands, Hastings  
sales@ahdltd.co.nz

[WWW.AHDLTD.CO.NZ](http://WWW.AHDLTD.CO.NZ)

