

WARNING: KEEP OUT OF REACH OF CHILDREN.
FOR ANIMAL TREATMENT ONLY

Ⓟ PARALOCK™

Abamectin Pour-on for Cattle

For the treatment and control of internal
and external parasites of cattle.

ACTIVE INGREDIENT:

Abamectin 10 mg/mL. Contains >941 g/L diethylene glycol monobutyl ether.



Supporting Kiwi Farmers



Ⓟ

5L

PARALOCK™

Abamectin Pour-on for Cattle

Read the entire label before use.

WARNING Harmful if swallowed. Causes serious eye irritation. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs through prolonged or repeated exposure.

Handling Precautions: Do not handle until all safety precautions have been read and understood. Wear protective gloves/protective clothing/eye protection. Avoid skin and eye contact. Do not breathe mist/vapour/spray. Avoid contact during pregnancy and while nursing. Keep closed when not in use. Wash hands and exposed skin thoroughly after handling and before meals and after use. Do not eat, drink or smoke when using this product.

First Aid: If swallowed do NOT induce vomiting. Rinse mouth. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention. If exposed or concerned or you feel unwell, get medical advice/attention. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

Environmental Protection: Very toxic to aquatic life with long lasting effects. Hazardous to soil organisms. Hazardous to terrestrial vertebrates. Hazardous to terrestrial invertebrates. Avoid release to the environment. Collect spillage.

Emergency Response: In case of spillage wear appropriate protective clothing. Prevent material from entering waterways and absorb spills with inert material and place in waste container. Wash area and absorb with further inert material.

Disposal: Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill or if appropriate recycle.

DOSAGE:

The dose rate for cattle is 500 µg abamectin per kg bodyweight, which is provided by 1 mL Paralock Abamectin Pour-On per 20 kg bodyweight, or according to the table.

Weight range	Dose volume	Dose per 5 L Pack
101-150 kg	7.5 mL	666
151-200 kg	10 mL	500
201-250 kg	12.5 mL	400
251-300 kg	15 mL	333
301-350 kg	17.5 mL	285
351-400 kg	20 mL	250
401-450 kg	22.5 mL	222
451-500 kg	25 mL	200
501-550 kg	27.5 mL	181
551-600 kg	30 mL	166
601-650 kg	32.5 mL	153

Heavier animals (e.g. mature bulls) should receive an additional 1 mL for each 20 kg over 650 kg.

INDICATION:

Paralock Abamectin Pour-On, when used at the recommended dose level of 500 micrograms abamectin per kg liveweight, effectively controls the following parasites of cattle: Gastrointestinal roundworms: adult and immature *Ostertagia* spp (including inhibited immatures), *Haemonchus* spp, *Trichostrongylus* spp, *Cooperia* spp, *Bunostomum* spp, *Oesophagostomum* spp, and adults of *Chabertia* spp, *Nematodirus* spp, *Trichuris* spp, and *Strongyloides* spp.

Lungworms: adult and immature *Dictyocaulus viviparus*.

Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress to the animal. The product other than as directed on the label.

Paralock Abamectin Pour-On contains 100 mg/mL abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family.

Registered to: Animal Health Direct Ltd
2 Tumu Way, Mangakino, Hastings 4175
06 873 5611 www.ahdtd.co.nz



Registered pursuant to the ACVM Act 1997: A012190
See www.foodsafety.govt.nz for registration conditions.
EPA Approval number HSR100759
See www.epa.govt.nz for approval conditions.

Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment. Dose rate to be based on heaviest cattle in group (bulls, cows, steers, calves etc). Do not under-dose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Do not use with standard pour-on equipment. Apply along the backline from shoulders to tail. Paralock Abamectin Pour-On is rainfast, even if rain falls 2 hours or more after treatment.

Contraindication

Do not use this product in calves under 4 months of age and under 100 kg liveweight. It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS:

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

Meat: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 35 days of the last treatment.

Storage: Store locked up below 25°C. Protect from sunlight and keep it well closed.



Batch:

Expiry:



ENVIRONMENTALLY HAZARDOUS SUBSTANCE,
LIQUID, N.O.S. (Contains Abamectin)
UN No 3082, Packing Group III, HazChem 3Z

