

AHD LTD PARALOCK™

TRIPLE SHEEP ORAL

Triple active combination oral drench for sheep.

NEW PRODUCT RELEASE



PRODUCT USAGE





For the treatment and control of all major gastrointestinal parasites in sheep, including strains with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families.

Also treats and controls lungworm, itch mite, tapeworm, larval stages of nasal bot, and aids in the control of adult liver fluke infection in sheep. Provides a supplemental source of copper, selenium, zinc, iodine and cobalt.

Refer to label for complete information and full chart of dose rates.

Paralock™ drenches will be presented in the AHD Ltd distinctive olive-green containers complete with drum tap located behind the back label.

PRODUCT BENEFITS

-  Reduced meat withholding period of 14 days. Milk 35 days.
-  Prolongs the effectiveness of all drench families.
-  Effective against worms for sheep with single and dual resistance.
-  Aids in the control of adult liver fluke.



PARALOCK INDICATIONS:

Sheep: For the effective treatment and control of mature and immature stages of susceptible strains of the following gastrointestinal parasites (* includes inhibited fourth stage larvae):

- Teladorsagia circumcincta*
- Teladorsagia trifurcata*
- Haemonchus contortus*
- Trichostrongylus axei
- Trichostrongylus colubriformis
- Trichostrongylus vitrinus
- Cooperia curticei*
- Cooperia oncophora*
- Cooperia punctata*
- Trichuris ovis
- Chabertia ovina
- Oesophagostomum venulosum
- Strongyloides papillosus
- Nematodirus filicollis
- Nematodirus spathiger.

Also effective against:

- Tapeworm (*Moniezia expansa*)
- Lungworm (*Dictyocaulus filaria*)
- Itch mite (*Psorobia ovis*)
- Nodule worm (*Oesophagostomum columbianum*) and all larval stages of nasal bot (*Oestrus ovis*)
- Aids in the control of adult liver fluke (*Fasciola hepatica*) infestations in sheep.

Anthelmintic resistant strains:

For the effective treatment and control of those gastrointestinal worms that have single and dual resistance to this combination of benzimidazole, levamisole/ morantel or avermectin/milbemycin anthelmintic action families.

Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

Ask your local parasitologist or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

DOSE RATES:

1mL per 5kg body weight equivalent to albendazole 5mg, levamisole hydrochloride 8mg, abamectin 0.2mg, selenium 0.1mg, copper 0.42mg, zinc 0.12mg, cobalt 0.04mg and iodine 0.15mg per kg body weight.

Body Weight	Dose	Doses per 20 L
20 kg	4.0 mL	5000
21-25 kg	5.0 mL	4000
26-30 kg	6.0 mL	3333
31-35 kg	7.0 mL	2857
36-40 kg	8.0 mL	2500
41-45 kg	9.0 mL	2222
46-50 kg	10.0 mL	2000
51-55 kg	11.0 mL	1818
56-60 kg	12.0 mL	1666
61-65 kg	13.0 mL	1538
66-70 kg	14.0 mL	1428
71-75 kg	15.0 mL	1333

Over 75 kg add 1mL for every 5kg body weight.

Drum tap located behind back label.

ACTIVE INGREDIENTS

Abamectin	1 g/L
Levamisole Hydrochloride	40 g/L
Albendazole	25 g/L
Copper	2.1 g/L
Selenium	0.5 g/L
Zinc	0.6 g/L
Cobalt	0.21 g/L
Iodine	0.76 g/L

CONTACT US

Animal Health Direct Ltd
2 Tumu Way, Longlands, Hastings
sales@ahdltd.co.nz

WWW.AHDLTD.CO.NZ

