

# AHD LTD PARALOCK™

## TRIPLE + TAPE ORAL

Triple active combination plus tape oral drench for sheep.

## NEW PRODUCT RELEASE






## PRODUCT USAGE

Paralock™ Triple Tape contains albendazole, a member of the benzimidazole family, levamisole, a member of the imidazothiazole family and abamectin, a member of the macrocyclic lactone family. It is effective against gastrointestinal worms sensitive to one, two or all three broad spectrum drench families. It also contains praziquantel, a member of the isoquinoline family, which is effective against tapeworm. Provides a supplemental source of selenium, copper, cobalt, zinc, and iodine.

Refer to label for complete information and full chart of dose rates.

Paralock™ drenches are presented in the AHD Ltd distinctive olive-green containers complete with drum tap located behind the back label.

## PRODUCT BENEFITS

-  Meat withholding period of 21 days.  
Milk 35 days.
-  Dosage 1mL / 5kg.
-  Effective against gastrointestinal worms that have single and dual resistance including tape worm.



## PARALOCK INDICATIONS:

**Sheep:** For the effective treatment and control of mature and immature stages of the following gastrointestinal parasites (\*includes inhibited fourth stage larvae):

- Teladorsagia circumcincta\*
- Teladorsagia trifurcata\*
- Haemonchus contortus\*
- Trichostrongylus axei
- Trichostrongylus colubriformis,
- Trichostrongylus vitrinus
- Cooperia curticei\*
- Cooperia oncophora\*
- Cooperia punctata\*
- Trichuris ovis
- Chabertia ovina
- Oesophagostomum venulosum
- Strongyloides papillosus
- Nematodirus filicollis and Nematodirus spathiger

Also effective against all stages of:

- Tapeworm (Moniezia expansa)
- Lungworm (Dictyocaulus filaria)
- Itch mite (Psorobia ovis)
- Nodule worm (Oesophagostomum columbianum)

And all larval stages of:

Nasal bot (Oestrus ovis) and as an aid in the control of adult liver fluke infestations in sheep.

### Anthelmintic resistant strains:

For the effective treatment and control of those gastrointestinal worms that have single and dual resistance to this combination of benzimidazole, levamisole/morantel or avermectin/milbemycin anthelmintic action families.

Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

## DOSE RATES:

1 mL per 5kg bodyweight equivalent to albendazole 5mg, levamisole hydrochloride 8mg, abamectin 0.2mg, praziquantel 3.8mg, selenium 0.1mg, copper 0.42mg, cobalt 0.04mg, zinc 0.12mg and iodine 0.15mg per kg bodyweight.

Bodyweight	Dose	10L treats (head)
20 kg	4 mL	2500
21-25 kg	5 mL	2000
26-30 kg	6 mL	1666
31-35 kg	7 mL	1428
36-40 kg	8 mL	1250
41-45 kg	9 mL	1111
46-50 kg	10 mL	1000
51-55 kg	11 mL	909
56-60 kg	12 mL	833
61-65 kg	13 mL	769
66-70 kg	14 mL	714
71-75 kg	15 mL	666

Over 75 kg add 1mL for every 5kg bodyweight.

## ACTIVE INGREDIENTS

Abamectin	1 g/L
Levamisole Hydrochloride	40 g/L
Albendazole	25 g/L
Praziquantel	19 g/L
Copper	2.1 g/L
<b>Selenium</b>	0.5 g/L
Zinc	0.6 g/L
Cobalt	0.21 g/L
Iodine	0.76 g/L

## CONTACT US

Animal Health Direct Ltd  
2 Tumu Way, Longlands, Hastings  
sales@ahdltd.co.nz

**WWW.AHDLTD.CO.NZ**

