

UNLOCK™

CYROMAZINE SPRAY-ON



For the residual protection of sheep against flystrike caused by all the commonly found strike flies including the Australian blowfly (*Lucilia cuprina*) for up to 8 weeks.

ACTIVE INGREDIENT:
60 g/L cyromazine



NET CONTENTS
20 Litres



WARNING: Dangerous to the environment. Harmful to aquatic life. Avoid contamination of any water supply with product or empty container.



PLEASE READ LABEL COMPLETELY BEFORE USE

HANDLING PRECAUTIONS: Causes skin irritation. Causes serious eye damage. Avoid contact with skin and eyes. Wear eye and face protection. Cyromazine may possibly affect blood and the haemopoietic system. Acetic acid possibly may damage lungs following inhalation. Do not breath fumes, vapours and spray. Handle with care. Wash hands and exposed skin before meals and after work.

FIRST AID: If medical advice is needed, have product container or label on hand. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before re-use. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. For advice, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately.

EMERGENCY RESPONSE: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

DISPOSAL: Preferably, dispose of product by use. Otherwise, dispose of product and packaging at an approved landfill or other approved facility. Triple rinse container and dispose of rinsate away from waterways. Burn in an appropriated incinerator, if circumstances such as wind direction permit, otherwise crush or punture and bury in a suitable landfill.

GENERAL INFORMATION: Unlock Cyromazine Spray-On is an easy use spray-on that will give residual protection against flystrike caused by all the commonly found strike flies including the Australian blowfly (*Lucilia cuprina*) for up to 8 weeks. Unlock Cyromazine Spray-On is to be used only on coarse woolled breeds of sheep. Unlock Cyromazine Spray-On is suitable for use on sheep between 4 and 12 weeks after shearing. The period of protection may be reduced by incorrect application; excessive fleece soiling; heavy fleece conditioning; if infections such as mycotic dermatitis (fleece rot) are present; or if prolonged heavy rain occurs following applicaton.

DIRECTIONS FOR USE - SHAKE GENTLY BEFORE USE

By law, the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress when using the product in a manner other than as recommended on the label.

Apply Unlock Cyromazine Spray-On through a suitable clean applicator gun. The product should be sprayed on in bands with the gun held 20cm above the wool during application. Unlock Cyromazine Spray-On should be applied onto those areas where maximum protection against flystrike is required.

Unlock Cyromazine Spray-On applied between the rump and shoulders should be sprayed on in even bands as shown (see diagram). The product sprayed onto the rump should be applied in an arc over the area shown (see diagram). The dose rate and number of bands to be applied are shown in the table. Note: The dose volume should be based on weight of the heaviest sheep in the mob.

Application and dose rate of Unlock Cyromazine Spray-On for the protection of all the commonly struck areas of the sheep (Table A).





Table A	2 bands for Sheep between 11-25kg eg. lambs			4 bands for sheep between 26-50kg eg. hoggets				5 bands for sheep between 51-70kg eg. ewes	
Liveweight (kg)	11-15	16-20	21-25	26-30	31-35	36-40	41-50	51-60	61-70
No. of bands	2	2	2	4	4	4	4	5	5
Vol. per band (mL)	10	14	18	11	13	15	17.5	17	20
Total vol applied (mL)	20	28	36	44	52	60	70	85	100
Sheep treated per 20L	1000	714	555	454	384	333	285	235	200

Application and dose rate of Unlock Cyromazine Spray-On for the protection of the crutch only (Table B).






Table B	2 bands for Sheep between 11-25kg eg. lambs			3 bands for sheep between 26-50kg eg. hoggets			4 bands for sheep between 51-70kg eg. ewes		
Liveweight (kg)	11-15	16-20	21-25	26-30	31-35	36-40	41-50	51-60	61-70
No. of bands	2	2	2	3	3	3	3	4	4
Vol. per band (mL)	7.5	10.5	13.5	11	13	15	17.5	16	18.5
Total vol applied (mL)	15	21	27	33	39	45	52.5	64	74
Sheep treated per 20L	1333	952	740	606	512	444	380	312	270

Application and dose rate of Unlock Cyromazine Spray-On at docking for prevention of flystrike. Apply between 5 & 10mL over the crutch ensuring that sufficient product to cover the area to be protected is applied.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

MEAT: Sheep producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

Wool Residues: Ensure the time between dipping and subsequent shearing is at least two months, except in the case of spot treatment of flystrike. For advice on your dipping programme, contact your supplier.

Conditions of Sale: Product sold by Animal Health Direct Ltd. will be suitable for use in the manner stated on the label. As Animal Health Direct Ltd. do not have any control over the way in which this chemical is used and the timing of application, Animal Health Direct Ltd. will not guarantee any application results. Liability will not be accepted for any loss arising through application, handling, storage and transportation of goods.

Storage: Store in original container, tightly closed, away from direct sunlight, in a cool, dry place, avoiding temperatures below 0° and above 35°C.

EPA approval number HSR100759.

See www.epa.govt.nz for approval conditions.

Registered Pursuant to the ACVM Act 1997.

No. A11884. See www.foodsafety.govt.nz

for registration conditions.

Registered to: Seneca Holdings Limited,

PO Box 58881, Botany, Auckland

Distributed by: Animal Health Direct Ltd.

1229 Maraekakaho Road, Hastings, New Zealand.

sales@ahdLtd.co.nz. Ph (06) 873 3611

Batch No.

Expiry date:

