

**WARNING - KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

AHD **UNLOCK**™

CYROMAZINE SPRAY-ON

Ready to use spray-on sheep dip
for the preventative control of
blowfly strike on sheep.

SHAKE WELL BEFORE USE

ACTIVE INGREDIENT:
60 g/L Cyromazine

NET CONTENTS
20 Litres



AHD UNLOCK™ CYROMAZINE SPRAY ON

PLEASE READ THIS LABEL COMPLETELY BEFORE USE

Environmental Protection Harmful to aquatic life. Avoid contamination of any water supply with product or empty container.

Handling Precautions: Causes skin irritation. Causes serious eye irritation. Avoid contact with skin and eyes. Do not breathe mist or spray. Wear protective gloves and eye and face protection. Cyromazine may possibly affect blood and the haemopoietic system. Handle with care. Wash hands and exposed skin before meals and after work.

First Aid: If skin or hair contact occurs, flush skin and hair with running water. Take off contaminated clothing and wash before re-use. If skin irritation occurs, get medical advice. If eye contact occurs, hold eyes open and rinse cautiously with water for several minutes. If you feel unwell or advice is required, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately.

Emergency Response: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

Disposal: Preferably, dispose of product by use. Otherwise, dispose of product and packaging at an approved landfill or other approved facility. Triple rinse container and dispose of rinsate away from waterways. Burn in an appropriate incinerator, if circumstances such as wind direction permit, otherwise crush or puncture and bury in a suitable landfill.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Meat: Sheep producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment.

Milk: Milk intended for sale for human consumption must be discarded during treatment for not less than 35 days following the last treatment.

WOOL RESIDUES: Ensure the time between dipping and subsequent shearing is at least two months, except in the case of spot treatment of flystrike. For advice on your dipping programme contact your supplier.

ACTIVE INGREDIENT: AHD UNLOCK CYROMAZINE SPRAY-ON contains 60g/litre cyromazine in the form of an aqueous concentrate.

GENERAL INFORMATION: AHD UNLOCK CYROMAZINE SPRAY-ON is an easy-to-use spray-on, which will give up to 6 weeks residual protection against flystrike caused by all the commonly found strike flies including the Australian blowfly (*Lucilia cuprina*). AHD UNLOCK CYROMAZINE SPRAY-ON can be used on all breeds of sheep. AHD UNLOCK CYROMAZINE SPRAY-ON is suitable for use on sheep between 4 and 12 weeks after shearing.

A shorter period of protection may occur if AHD UNLOCK CYROMAZINE SPRAY-ON is applied incorrectly, if there is excessive fleece soiling or if wool infections such as mycotic dermatitis (fleece rot) are present. Prolonged heavy rain following treatment may also reduce the protection period provided by AHD UNLOCK CYROMAZINE SPRAY-ON.





DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Apply AHD UNLOCK CYROMAZINE SPRAY-ON through a suitable clean applicator gun. The product should be sprayed on in bands with the gun held 20cm above the wool during

application. AHD UNLOCK CYROMAZINE SPRAY-ON should be applied onto those areas where maximum protection against flystrike is required. AHD UNLOCK CYROMAZINE SPRAY-ON applied between the rump and shoulders should be sprayed on in even bands as shown (see diagram). The product sprayed onto the rump should be applied in an arc over the area shown (see diagram). The dose rate and number of bands to be applied are shown in the table.

Note: The dose volume applied should be based on the weight of the heaviest sheep in the mob.

APPLICATION AND DOSE RATE OF AHD UNLOCK CYROMAZINE SPRAY-ON FOR THE PROTECTION OF ALL THE COMMONLY STRUCK AREAS OF THE SHEEP

							
	2 bands Sheep between 11-20kg	3 bands Sheep between 21-40kg	4 bands Sheep between 41-60kg	5 bands Sheep between 61-70kg			
Liveweight kg	11-15	16-20	21-30	31-40	41-50	51-60	61-70
No. of bands	2	2	3	3	4	4	5
Volume per band (mL)	7.5	10	10	13.3	12.5	15	14
Total vol applied (mL)	15	20	30	40	50	60	70
Sheep treated per 20L	1333	1000	666	500	400	333	285



Application and dose rate of AHD UNLOCK CYROMAZINE SPRAY-ON at docking for the prevention of flystrike Apply between 5mL and 10mL over the crutch ensuring that sufficient product to cover the area to be protected is applied (see diagram).

STORAGE: Store in original container, tightly closed, away from direct sunlight, in a cool, dry place, avoiding temperatures below 0°C and above 35°C.

DISPOSAL OF CONTAINER: Do not reuse empty container but burn or flatten and bury in a suitable landfill or if appropriate recycle.

Conditions of Sale: Product sold by Animal Health Direct Ltd. will be suitable for use in the manner stated on the label. As Animal Health Direct Ltd. do not have any control over the way in which this chemical is used and the timing of application, Animal Health Direct Ltd. will not guarantee any application results. Liability will not be accepted for any loss arising through application, handling, storage and transportation of goods.

Registered pursuant to the ACVM Act 1997, No. A11946
See www.foodsafety.govt.nz for registration conditions.

Registered to: WSD Agribusiness Pty Ltd
Distributed by: Animal Health Direct Ltd.
1229 Maraekakaho Road, Hastings, New Zealand.
sales@ahdLtd.co.nz Ph (06) 873 3611.

Batch No.

Expiry date:

